

Product Information: Jevity® 1.2 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Jevity® 1.2 Cal

Complete, Balanced Nutrition® with Fiber

- 1 JEVITY 1.2 CAL is fiber-fortified therapeutic nutrition that provides complete, balanced nutrition for long- or short-term tube feeding.
- 1 For tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 May be used for oral feeding of patients with altered taste perception.
- 1 Use under medical supervision.
- 1 Concentrated calories (1.2 Cal/mL) to help tube-fed patients gain and maintain healthy weight.
- 1 4 g dietary fiber/8 fl oz (2.1 g of a unique fiber blend and 1.9 g of scFOS) which helps meet daily fiber intake. Fiber helps moderate bowel function.^{1,2,3}
- 1 Includes 1.9 g of scFOS/8 fl oz (8 g/L and 12 g/1.5L). scFOS are prebiotics that stimulate the growth of beneficial bacteria in the colon.^{1,2,3}
- 1 Meets or exceeds 100% of the RDIs for 25 essential vitamins and minerals in 1500 Cal (1250 mL).
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not for IV use.
- 1 Not suitable for people with galactosemia.

¹ *Bornet FR, et al. Nutr Rev. 2002;60(11):326-334.*

² *Tokunaga T, et al. Bifidus. 1993;6(2):143-150.*

³ *Bouhnik Y, et al. Nutr J. 2007;6:42-48.*

Ingredients

Liquid Unflavored:

Water, Corn Maltodextrin, Corn Syrup Solids, Sodium Caseinate, Soy Protein Isolate, Canola Oil, Corn Oil, Short-chain Fructooligosaccharides, Medium Chain Triglycerides, Calcium Caseinate, Soy Fiber. Less than 0.5% of the Following: Oat Fiber, Potassium Citrate, Calcium Phosphate, Magnesium Phosphate, Gum Arabic, Potassium Chloride, Soy Lecithin, Cellulose Gum, Choline Chloride, Ascorbic Acid, Salt, Potassium Hydroxide, L-Carnitine, Taurine, dl-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Sodium Citrate, Zinc Sulfate, Manganese Sulfate, Niacinamide, Copper Sulfate, Calcium Pantothenate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Beta-Carotene, Vitamin A Palmitate, Chromium Chloride, Folic Acid, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Biotin, Phyloquinone, Vitamin D3, and Vitamin B12.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
64625	Jevity 1.2 Cal / 8 fl oz (237 mL) Recloseable Carton / 24 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)	
	Value	%RDI / %DV
Calories	285	
Protein, g	13.2	
Fat, g	9.3	
Carbohydrate, g	40.2	
Dietary Fiber, g	4*	
Total Sugars, g	3	
Added Sugars, g	3	
L-Carnitine, mg	36	
Taurine, mg	36	
Water, mL	191	
Vitamin A, mcg RAE	171†	
Beta-Carotene, mcg	307	
Vitamin D, mcg	3.8	
Vitamin E, mg	9.2	
Vitamin K, mcg	29	
Vitamin C, mg	59	
Folate, mcg DFE	76	
Folic Acid, mcg	46	
Thiamin (Vitamin B1), mg	0.41	
Riboflavin (Vitamin B2), mg	0.59	
Vitamin B6, mg	0.33	
Vitamin B12, mcg	2.3	
Niacin, mg NE	4	
Choline, mg	142	
Biotin, mcg	5.7	
Pantothenic Acid, mg	0.95	
Sodium, mg	253	
Potassium, mg	566	
Chloride, mg	356	
Calcium, mg	284	
Phosphorus, mg	284	
Magnesium, mg	88	
Iodine, mcg	36	
Manganese, mg	0.81	
Copper, mg	0.41	
Zinc, mg	2.5	
Iron, mg	3.4	
Selenium, mcg	14	
Chromium, mcg	14	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)	
	Value	%RDI / %DV
Molybdenum, mcg	23	

Liquid Unflavored Footnotes & References

Per 8 fl oz (237 mL)

* 2.1 g of total dietary fiber from a unique fiber blend and 1.9 g of short-chain fructooligosaccharides.

† 122 mcg RAE of vitamin A activity supplied by 307 mcg of beta-carotene.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Instructions for Use:

8 fl oz Container

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Shake well prior to opening.
- 1 After opening, refrigerate and consume product within 48 hours. If consuming product directly from the container, refrigerate and consume within 24 hours.

Oral Feeding

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

Tube Feeding

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Feed at room temperature using a feeding pump or syringe.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.

Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **INSTRUCTIONS FOR USE** increases the potential for microbial contamination and may reduce hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from RTH Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.

Precautions

- 1 Follow directions for use provided by manufacturer of feeding sets.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.